

OTC use in Norway for ulipristal acetate, ATC-code: G03AD02

- This OTC substance report is based on the assessment of the OTC indication and posology for products containing ulipristal acetate. It defines the preferred Norwegian wording for the package leaflet and labelling for OTC products containing ulipristal acetate. In addition, an overview of the approvable strength(s), pharmaceutical form(s) and pack size(s) exempt from medical prescription in Norway is included.
- The proposed OTC indication and posology in the OTC package leaflet and labelling must be covered by the information approved in the corresponding SmPC.

Preparations for oral use, up to 30mg per unit

1. Package leaflet

1.1 Indication

Til Kvinner: nødprevensjon etter ubeskyttet samleie eller prevensjonssvikt.

1.2 Posology

Change the quantity from the given strength to the number of entities to be taken (e.g. 1-2 tablets, 1 suppository, 20 ml...).

Kvinner: ta 30 mg så snart som mulig og ikke senere enn 120 timer (5 døgn) etter ubeskyttet samleie eller prevensjonssvikt.

2. Labelling

2.1 Indication

State the indication as in the PIL. If the full indication is stated on the back panel of the package, the following abbreviation can be used on the front panel:

Nødprevensjon

2.2 Posology

State the dosage as in the PIL.

2.3 Other information

Not applicable.

3. Content of the pack

The table below presents the highest level of the terms for approvable pharmaceutical forms, if possible. For example: the term “tablets” includes all types of tablet formulations as for example film coated tablets or chewable tablets. For active substances where some pharmaceutical forms are exempt from approval due to safety concern, this is stated explicitly below the table.

Pharmaceutical form	Maximum strength	Maximum pack size
Tablet	30 mg	1